

## Examen de Matemáticas 2ºBachillerato(CN) Enero 2008

---

**Problema 1** Calcular los siguientes límites:

- a)  $\lim_{x \rightarrow 0} \frac{x - \sin x}{\tan x - \sin x}$  (Castilla La Mancha 2005)
- b) Si  $f(x) = x^3 e^{-x}$  calcular  $\lim_{x \rightarrow \infty} f(x)$  y  $\lim_{x \rightarrow -\infty} f(x)$  (Isla Baleares 2005)
- c)  $\lim_{x \rightarrow 3} \frac{\cos(\pi x)}{x}$  (La Rioja 2005)
- d)  $\lim_{x \rightarrow \infty} (\sqrt{x^2 + x} - \sqrt{x^2 - x})$  (Madrid 2005)
- e)  $\lim_{x \rightarrow \infty} x \left[ \arctan(e^x) - \frac{\pi}{2} \right]$  (Madrid 2005)
- f)  $\lim_{x \rightarrow 0} \frac{4x + \sin 2x}{\sin 3x}$  (Zaragoza)
- g)  $\lim_{x \rightarrow \infty} \left( \frac{2x - 1}{2x} \right)^{x+2}$
- h)  $\lim_{x \rightarrow 0} x^x$
- i)  $\lim_{x \rightarrow \infty} \frac{\sqrt{3x^2 + x - 1}}{2x + 3}$
- j)  $\lim_{x \rightarrow 0} \frac{\sqrt{3x + 1} - \sqrt{x + 1}}{x}$

**Solución:**

a)  $\lim_{x \rightarrow 0} \frac{x - \sin x}{\tan x - \sin x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \lim_{x \rightarrow 0} \frac{1 - \cos x}{1 + \tan^2 x - \cos x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} =$   
 $\lim_{x \rightarrow 0} \frac{\sin x}{2 \tan x (1 + \tan^2 x) + \sin x} = \lim_{x \rightarrow 0} \frac{\sin x}{2 \tan x + 2 \tan^3 x + \sin x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} =$   
 $\lim_{x \rightarrow 0} \frac{\cos x}{2(1 + \tan^2 x) + 6 \tan^2 x (1 + \tan^2 x) + \cos x} = \frac{1}{3}$

b)  $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} x^3 e^{-x} = [\infty \cdot 0] = \lim_{x \rightarrow \infty} \frac{x^3}{e^x} = \begin{bmatrix} \infty \\ \infty \end{bmatrix} = \lim_{x \rightarrow \infty} \frac{6x}{e^x} = \begin{bmatrix} \infty \\ \infty \end{bmatrix} = \lim_{x \rightarrow \infty} \frac{6}{e^x} = 0$

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow -\infty} x^3 e^{-x} = \lim_{x \rightarrow -\infty} -x^3 e^x = -\infty$$

c)  $\lim_{x \rightarrow 3} \frac{\cos(\pi x)}{x} = -\frac{1}{3}$

d)

$$\begin{aligned} & \lim_{x \rightarrow \infty} (\sqrt{x^2 + x} - \sqrt{x^2 - x}) = [\infty - \infty] = \\ & \lim_{x \rightarrow \infty} \frac{(\sqrt{x^2 + x} - \sqrt{x^2 - x})(\sqrt{x^2 + x} + \sqrt{x^2 - x})}{\sqrt{x^2 + x} + \sqrt{x^2 - x}} = \\ & \lim_{x \rightarrow \infty} \frac{(\sqrt{x^2 + x})^2 - (\sqrt{x^2 - x})^2}{\sqrt{x^2 + x} + \sqrt{x^2 - x}} = \lim_{x \rightarrow \infty} \frac{2x}{\sqrt{x^2 + x} + \sqrt{x^2 - x}} = \\ & \lim_{x \rightarrow \infty} \frac{\frac{2x}{x}}{\sqrt{\frac{x^2 + x}{x^2}} + \sqrt{\frac{x^2 - x}{x^2}}} = \frac{2}{2} = 1 \end{aligned}$$

e)

$$\begin{aligned} & \lim_{x \rightarrow \infty} x \left[ \arctan(e^x) - \frac{\pi}{2} \right] = [0 \cdot \infty] = \lim_{x \rightarrow \infty} \frac{\arctan(e^x) - \frac{\pi}{2}}{1/x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \\ & \lim_{x \rightarrow \infty} \frac{\frac{e^x}{1+e^{2x}}}{-\frac{1}{x^2}} = \lim_{x \rightarrow \infty} \frac{-x^2 e^x}{1+e^{2x}} = \begin{bmatrix} \infty \\ \infty \end{bmatrix} = \lim_{x \rightarrow \infty} \frac{-2xe^x - x^2 e^x}{2e^{2x}} = \\ & \lim_{x \rightarrow \infty} \frac{-2x - x^2}{2e^x} = \begin{bmatrix} \infty \\ \infty \end{bmatrix} = \lim_{x \rightarrow \infty} \frac{-2 - 2x}{2e^x} = \begin{bmatrix} \infty \\ \infty \end{bmatrix} = \lim_{x \rightarrow \infty} \frac{-2}{2e^x} = 0 \end{aligned}$$

f)  $\lim_{x \rightarrow 0} \frac{4x + \sin 2x}{\sin 3x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \lim_{x \rightarrow 0} \frac{4 + 2 \cos 2x}{3 \cos 3x} = \frac{6}{3} = 2$

g)  $\lim_{x \rightarrow \infty} \left( \frac{2x - 1}{2x} \right)^{x+2} = e^{-1/2}$

h)  $\lim_{x \rightarrow 0} x^x = \lambda \implies \lim_{x \rightarrow 0} x \ln x = \ln \lambda$

$$\lim_{x \rightarrow 0} x \ln x = \lim_{x \rightarrow 0} \frac{\ln x}{x} = \begin{bmatrix} -\infty \\ \pm\infty \end{bmatrix} = \lim_{x \rightarrow 0} \frac{1/x}{-1/x^2} = \lim_{x \rightarrow 0} (-x) = 0$$

$$\ln \lambda = 0 \implies \lambda = 1$$

i)  $\lim_{x \rightarrow \infty} \frac{\sqrt{3x^2 + x - 1}}{2x + 3} = \frac{\sqrt{3}}{2}$

j)  $\lim_{x \rightarrow 0} \frac{\sqrt{3x + 1} - \sqrt{x + 1}}{x} = 1$